

JK



Office de la Propriété  
Intellectuelle  
du Canada  
Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office  
An agency of  
Industry Canada

CA 2034096 (2002/03/26)

(11)(21) 2 034 096

(12) BREVET CANADIEN  
CANADIAN PATENT

(13) C

(2) Date de dépôt / filing Date: 1991/01/14

(41) Mise à la disp. pub. / Open to Public Insp.: 1991/07/16

(42) Date de délivrance / Issue Date: 2002/03/26

(43) Priorité / Priority: 1990/01/15 (149790) II

(51) Cl. Int. / Int. Cl. 5 A61K 31/19, A61K 9/24

(62) Inventeurs / Inventors:  
Bourke, Edward Anthony, IE;  
Mulligan, Seamus, II(73) Propriétaire / Owner:  
ELAN CORPORATION P.C., II

(74) Agent: KIRBY EADES GALE BAKER

(51) Titre: FORMULATION DE NAPROXENE A LIBERATION CONTROLÉE POUR ADMINISTRATION UNE FOIS PAR JOUR

(51) Title: CONTROLLED ABSORPTION NAPROXEN FORMULATION FOR ONCE-DAILY ADMINISTRATION

## (57) Abrégé / Abstract

A naproxen formulation for oral administration on a once-daily basis comprises a first portion of the naproxen formulated in a multi-particulate pellet form designed to release the drug at a rate sufficient to maintain therapeutically effective blood levels substantially over 24 hours when administered each day on a once-daily basis and, optionally, a second portion of the naproxen formulated so as to release the drug promptly following administration so as to obtain a relatively immediate therapeutic response.

Canada

<http://cipo.gc.ca> • Ottawa-Hull K1A 0G9 • [669-9696](http://cipo.gc.ca)  
CIPC • CIPU 101

O P I C



C I P O

2034096

This invention relates to a controlled absorption pharmaceutical formulation and, in particular, to a controlled absorption form of naproxen for oral administration.

5 Naproxen, a propionic acid derivative ((S) - 6-methoxymethyl-2-naphthalenecarboxylic acid), is a non-steroidal, anti-inflammatory drug (NSAID) which exhibits analgesic and antipyretic properties. The exact mechanisms of action have not been clearly established, but many of the effects of naproxen are associated with the inhibition of prostaglandin synthesis and in particular cyclo-oxygenase, an enzyme that catalyzes the formation of prostaglandin precursors from arachidonic acid. Naproxen  
10 is used to relieve mild to moderately severe pain in rheumatoid arthritis, osteoarthritis and other inflammatory complaints.

Naproxen has been available for use for over a decade and has been found to be acceptably non-toxic by many regulatory agencies.  
15 Naproxen is used as either its free acid or its sodium salt, naproxen sodium. The present application concerns a naproxen sodium formulation, however, it will be understood that naproxen free acid could also be used.

Naproxen and naproxen sodium are conventionally administered on a two to four times daily basis. Plasma naproxen concentrations of 30-90 µg/ml reportedly are required for anti-inflammatory or analgesic effects. Reduced pain intensity has been demonstrated in sixty postpartum women from 0.5 to 6 hours after oral administration of naproxen or its sodium salt in doses sufficient to yield plasma naproxen  
20 levels between 30-70 µg/ml (Sevelius, H. *et al.*, (1980); Br. J. Clin. Pharmacol. 10, 259-263). Evidence from twenty-four patients with rheumatoid arthritis suggested that clinical response occurred at plasma naproxen levels above 50 µg/ml. (Day, R.O. *et al.*, (1982); Clin. Pharmacol. Ther. 31, 733-740). Thus, while the rate of absorption may  
25 affect the onset of analgesic activity, continued plasma levels of the drug  
30 are likely to be important in maintaining analgesia.

CA 02034096 2001-01-19

2

It is an object of the present invention to provide a controlled absorption naproxen formulation suitable for once-daily administration, and particularly one which demonstrates an initial rate of absorption resulting in a rapid analgesic effect and then exhibits controlled absorption characteristics resulting in continued plasma levels of the drug over a 24 hour period.

Accordingly, the invention provides a naproxen formulation for oral administration on a once-daily basis, said formulation comprising a first portion of the naproxen formulated in a multi-particulate pellet form, each pellet having a core of naproxen or a pharmaceutically acceptable salt thereof surrounded by a multi-layer membrane, said membrane containing a major proportion of a pharmaceutically acceptable film-forming, water insoluble polymer and optionally a minor proportion of a pharmaceutically acceptable film-forming, water soluble polymer, said multi-particulate form being effective to release the drug at such a rate as to maintain therapeutically effective blood levels substantially over 24 hours when administered each day on a once-daily basis and, optionally, a second portion of said naproxen formulated so as to release the drug promptly following administration so as to obtain a relatively immediate therapeutic response.

Preferably said first portion is in pellet form comprising a core of naproxen or a pharmaceutically acceptable salt thereof in association with an organic acid, the naproxen component and the organic acid being present in a ratio of from 20:1 to 1:1, and a multi-layer membrane surrounding said core and containing a major

CA 02034096 2001-01-19

2 a

proportion of a pharmaceutically acceptable film-forming, water insoluble polymer and optionally a minor proportion of a pharmaceutically acceptable film-forming, water soluble polymer, the number of layers in said membrane and the ratio of said water soluble to water insoluble polymer being effective to permit release of said naproxen from said pellet at a rate allowing controlled absorption thereof at therapeutically effective blood levels over a 24 hour period following each daily oral administration. Preferred formulations of the present invention are those wherein the dissolution rate of said pellet, when measured *in vitro* in a type 1 dissolution basket apparatus according to U.S. Pharmacopoeia XXII in phosphate buffer at pH 7.2 and at 75 r.p.m. substantially corresponds to the following dissolution pattern:

- a) from 0 to 50% of the total naproxen is released after 1 hour of measurement in said apparatus;
- b) from 20 to 70% of the total naproxen is released after 2 hours of measurement in said apparatus; and
- 5 c) not less than 50% of the total naproxen is released after 4 hours of measurement in said apparatus.

When the dissolution rate of said pellet is measured *in vitro* in a type 2 dissolution basket apparatus according to U.S. Pharmacopoeia XXII in phosphate buffer at pH 7.4 and at 50 r.p.m. the dissolution rate  
10 substantially corresponds to the following dissolution pattern:

- a) from 20 to 70% of the total naproxen is released after 1 hour of measurement in said apparatus;
- b) not less than 50% of the total naproxen is released after 2 hours of measurement in said apparatus; and
- 15 c) not less than 75% of the total naproxen is released after 4 hours of measurement in said apparatus.

The invention also provides a controlled absorption naproxen formulation for oral administration, comprising pellets as hereinbefore defined, said formulation including a sufficient quantity of a rapid  
20 release form of naproxen to ensure prompt achievement of analgesically effective blood levels together with the prolonged effects described above. Such formulations comprising rapid and prolonged release components preferably have a dissolution rate which when measured in  
25 a type 1 dissolution basket apparatus according to U.S. Pharmacopoeia XXII in phosphate buffer at pH 7.2 and at 75 r.p.m. substantially corresponds to the following dissolution pattern:

- a) from 15 to 50% of the total naproxen is released after 0.5 hours of measurement in said apparatus;

- b) from 25 to 75% of the total naproxen is released after 1 hour of measurement in said apparatus; and
- c) not less than 65% of the total naproxen is released after 4 hours of measurement in said apparatus.

5        When the dissolution rate of said pellet is measured *in vitro* in a type 2 dissolution paddle apparatus according to U.S. Pharmacopocia XXII in phosphate buffer at pH 7.4 and at 50 r.p.m. the dissolution rate substantially corresponds to the following dissolution pattern:

- 10            a) from 25 to 60% of the total naproxen is released after 0.5 hours of measurement in said apparatus;
- b) from 35 to 75% of the total naproxen is released after 1 hour of measurement in said apparatus; and
- c) not less than 65% of the total naproxen is released after 4 hours of measurement in said apparatus.

15        The applicants have found in the case of the above formulations of the present invention, that therapeutically effective blood levels can be maintained substantially over 24 hours with peak plasma levels occurring between 2 and 16 hours, preferably between 4 and 10 hours. The present invention defines this desired time to peak plasma level as  
20        the T<sub>max</sub> of the formulation.

             Advantageously, the formulation comprises a blend of pellets as hereinbefore defined together with up to 60% by weight of said rapid release form of naproxen, preferably 20-45%.

25        Most preferably, the rapid release form of naproxen comprises a rapid release granulate.

The naproxen used in the preferred dosage forms of the present invention is in the form of a pharmaceutically acceptable salt thereof, especially the sodium salt thereof.

- 5 The organic acid is preferably represented by one or more of the following acids: adipic acid, ascorbic acid, citric acid, fumaric acid, malic acid, succinic acid or tartaric acid. Especially preferred acids are fumaric acid and citric acid.

- 10 The core also optionally contains a lubricant which is represented by one or more of the following: sodium stearate, magnesium stearate, stearic acid or talc. The naproxen and lubricant are preferably present in a ratio of from 10:1 to 200:1.

- 15 Preferably, the core comprises naproxen or a pharmaceutically acceptable salt thereof and the associated organic acid embedded in a polymeric material. The polymeric material may be rapidly soluble in water or insoluble in water or freely permeable to naproxen and water.

The term water soluble polymer as used herein includes polymers which are freely permeable to water, whilst the term water insoluble polymer as used herein includes polymers which are slightly permeable to water.

- 20 The polymeric material preferably consists solely of a water insoluble polymer or a polymer which is slightly permeable to water and aqueous solutions of naproxen. Alternatively, the polymeric material may consist solely of a water soluble polymer or a polymer which is freely permeable to aqueous solutions of naproxen and water.
- 25 The polymeric material of the core may include a combination of a water insoluble polymer with a water soluble polymer. The ratio of water soluble/freely permeable to water insoluble/slightly permeable polymer is determined by the particular combination of polymers selected.

The water soluble polymer is suitably polyvinyl alcohol, polyvinylpyrrolidone, methyl cellulose, hydroxypropyl cellulose, agar, carrageenan, xanthan, hydroxypropylmethyl cellulose or polyethylene glycol or a mixture thereof. An especially preferred water soluble  
5 polymer is polyvinylpyrrolidone.

A suitable polymer which is freely permeable to naproxen and water is a polymer sold under the Trade Mark EUDRAGIT RL.

The water insoluble polymer of the core is suitably ethyl cellulose, cellulose acetate, cellulose propionate (lower, medium or  
10 higher molecular weight), cellulose acetate propionate, cellulose acetate butyrate, cellulose acetate phthalate, cellulose triacetate, poly(methyl methacrylate), poly(ethyl methacrylate), poly(butyl methacrylate), poly(isobutyl methacrylate), poly(hexyl methacrylate), poly(isodecyl methacrylate), poly(lauryl methacrylate), poly(phenyl methacrylate),  
15 poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), poly(octadecyl acrylate), poly(ethylene), poly(ethylene) low density, poly(ethylene) high density, poly(propylene), poly(ethylene oxide), poly(ethylene terephthalate), poly(vinyl isobutyl ether), poly(vinyl acetate), poly(vinyl chloride) or polyurethane or a mixture thereof.

20 The water insoluble polymer of the core may also comprise naturally occurring polymers or resins. Especially suitable water insoluble, naturally occurring polymers include shellac, chitosan, gumjuniper or a mixture thereof.

A suitable polymer which is slightly permeable to naproxen and  
25 water is a polymer sold under the Trade Mark EUDRAGIT RS or a polymer whose permeability is pH dependent and sold under the Trade Mark EUDRAGIT L, EUDRAGIT S or EUDRAGIT E. Especially preferred polymers in this category are EUDRAGIT L.

EUDRAGIT polymers are polymeric lacquer substances based on  
30 acrylates and/or methacrylates. The polymeric materials sold under the Trade Mark EUDRAGIT RL and EUDRAGIT RS are acrylic resins



- comprising copolymers of acrylic and methacrylic acid esters with a low content of quaternary ammonium groups and are described in the "EUDRAGIT" brochure of Messrs. Rohm Pharma GmbH (1985) wherein detailed physical-chemical data of these products is given. The ammonium groups are present as salts and give rise to the permeability of the lacquer films. EUDRAGIT RL and RS are freely permeable (RL) or slightly permeable (RS), respectively, independent of pH.

- EUDRAGIT L is an anionic polymer synthesized from methacrylic acid and methacrylic acid methyl ester. It is insoluble in acids and pure water. It becomes soluble in a neutral to weakly alkaline milieu by forming salts with alkalis. The permeability of EUDRAGIT L is pH dependent. Above pH 5.0, the polymer becomes increasingly permeable. EUDRAGIT L is described in the "EUDRAGIT L" brochure of Messrs. Rohm Pharma GmbH (1986) wherein detailed physical-chemical data of the product is given.

The core suitably has a number of layers of the core-forming materials and is built up in a manner known *per se*.

- A multi-layer arrangement of naproxen, organic acid and polymeric material is preferably built up on a central active core. The active core is formed by blending naproxen, organic acid and polymeric material to form a homogenous powder. A portion of the above blend is shaped to form a central core. A multi-layer arrangement is then built up by a successive layering and binding process wherein the remainder of the blend and a polymer binding solution are applied to the active core in alternative layers in a conventional coating pan. Alternatively, an automatic coating system may be used where the remainder of the blend and polymer binding solution is applied to the active core, simultaneously. Conventional automated coating systems include, for example, a CF granulator or a Glatt fluidized bed. The cores are formed to assure a uniform distribution of naproxen and excipient ingredients throughout the core. The preferred average diameter of the completed cores is in the range of 0.4 - 1.6 mm, an especially preferred average diameter being in the range of 0.6 - 1.2 mm.

The multi-layer arrangement of naproxen, organic acid and polymeric material may also be built up on a central inert core, suitably consisting of a non-pareil bead or seed of sugar/starch having an average diameter in the range 0.2-1.4 mm, especially 0.3-0.8 mm. The  
5 naproxen, organic acid and polymeric material may be built up on a central inert core as hereinbefore defined in a conventional coating pan or any automated coating system.

The naproxen, organic acid and optional other components are blended to form a homogenous powder. The naproxen component and  
10 organic acid component are preferably present in a ratio of from 20:1 to 1: 2, more especially 6:1 to 1:1. The blend is suitably passed through the appropriate mesh screen using a milling machine. In the case of coating in a conventional coating pan, alternate layers of a coating solution/suspension of the polymeric material and the powder are  
15 applied to the central inert core to build up the multi-layer arrangement of the core. In the case of an automatic coating system, the coating solution/suspension of the polymeric material and the powder are applied, simultaneously, in conventional manner. The coating solution/suspension of the polymeric material comprises one or more  
20 polymers dissolved/suspended in a suitable solvent or mixture of solvents. The concentration of the polymeric material in the coating solution/suspension is determined by the viscosity of the final solution/suspension. Preferably, between 5 and 60 parts of the central inert cores are used relative to the homogenous powder. The addition of  
25 a plasticizing agent to the polymeric solution/suspension may be necessary depending on the formulation to improve the elasticity and also the stability of the polymer film and to prevent changes in the polymer permeability over prolonged storage. Such changes could affect the drug release rate. Suitable plasticizing agents include  
30 polyethylene glycol, propylene glycol, glycerol, triacetin, dimethyl phthalate, diethyl phthalate, dibutyl phthalate, dibutyl sebacate, triethyl citrate, tributyl citrate, triethyl acetyl citrate, castor oil and varying percentages of acetylated monoglycerides.

As mentioned above, the core may optionally contain a lubricant. A preferred range of naproxen to lubricant ratio when a central inert core is used is 50:1 to 5:1.

5 Preferred coating materials include - solutions/suspensions of the polymers cited for use in the application of the powder blend to the central cores in a suitable organic/aqueous carrier medium.

10 The membrane of the film-forming polymer or mixture of polymers surrounding the core preferably has a proportion of a polymer which is slightly permeable to naproxen and water and optionally a proportion of a water permeable polymer, the ratio of slightly water permeable to water permeable polymer being determined by the inherent permeability characteristics of the polymer(s) selected.

15 The membrane may also be composed of a proportion of a polymer which is water insoluble and a proportion of a polymer which is water soluble, the ratio of water insoluble to water soluble polymer being determined by the inherent permeability characteristics of the respective polymers.

20 Normally the ratio of water insoluble/slightly permeable polymers to water soluble/permeable polymers lies between 1:5 and 50:1, more usually 1:2 and 20:1. Examples of each of these types of polymer are described above. Especially useful water soluble/permeable polymers include polyvinylpyrrolidone, polyvinyl alcohol and EUDRAGIT RL, whilst useful water insoluble/slightly permeable  
25 polymers include ethyl cellulose, cellulose acetate, EUDRAGIT RS, EUDRAGIT L, EUDRAGIT E and EUDRAGIT S. Commercially available ready-made polymeric solutions/suspensions may also be useful. These ready-made solutions/suspensions may optionally contain plasticizing agents to improve the polymer film as described previously. Examples of ready-made solutions/suspensions of polymeric material  
30 with or without plasticizing agent include EUDRAGIT RL 30D, EUDRAGIT NE 30D, EUDRAGIT E 12.5, EUDRAGIT L 12.5 P, EUDRAGIT E 12.5, EUDRAGIT S 12.5 P, EUDRAGIT RL 12.5,

10

2034096

EUDRAGIT RS 300, EUDRAGIT RS 12.5, AQUACOAT (a Trade Mark of PMC Corporation) and SURE-LEASE (a Trade Mark of Colorcon Inc.).

- 5 The water insoluble polymer of the membrane may also comprise naturally occurring polymers or resins. Especially suitable water insoluble, naturally occurring polymers include shellac, chitosan, gumjuniper or a mixture thereof.

- 10 The membrane may be built up by applying a plurality of coats of membrane polymer solution or suspension to the core as hereinafter described. The membrane solution or suspension contains the polymer(s) dissolved or suspended, respectively, in a suitable aqueous or organic solvent or mixture of solvents, optionally in the presence of a lubricant. Suitable lubricants are talc, stearic acid, magnesium stearate and sodium stearate. A particularly preferred lubricant is talc. The  
15 membrane, polymer or mixture of polymers may optionally include a plasticizing agent, the function and choice of which has been previously described.

The dissolution rate achieved is proportionally slower as the amount of membrane applied is increased.

- 20 The membrane solution or suspension may be applied to the active cores in a conventional coating pan as indicated or, alternatively, using an automated system such as a CF granulator, for example, a FREUND CF granulator, a GLATT fluidized bed processor, an AEROMATIC, a modified ACCELA-COTA or any other suitably  
25 automated bead coating equipment (FREUND, GLATT, AEROMATIC and ACCELA-COTA are all Trade Marks).

- 30 Preferably 2-75 ml of membrane solution/suspension is applied per application per kilogram of cores. In an automated system the total amount of membrane solution/suspension applied to the cores is the same as that applied in a conventional coating pan, except that the membrane solution/suspension may be applied continuously.

Preferably, when a coating pan is used the membrane is applied at a rate of 5-30 applications/day until all of the applications have been applied. Between days the pellets are dried for a suitable period of time at a controlled temperature.

- 5       The pellets and granulate may be compressed into tablets using a binder and/or hardening agent commonly employed in tableting such as microcrystalline cellulose sold under the Trade Mark "AVICEL" or a co-crystallized powder of highly modified dextrans (3% by weight) and sucrose sold under the Trade Mark "DI-PAC" in such a way that the  
10       specific dissolution rate of the pellets is maintained.

The pellets by themselves or combined with naproxen powder or naproxen cores may also be filled into hard or soft gelatine capsules.

- 15       Thus the applicants have incorporated multi-particulate pellets of the sodium salt of naproxen into a tablet base. These multi-particulate pellets are formulated with sustained release characteristics allowing a continuous supply of naproxen to be maintained in the plasma to achieve a prolonged analgesic effect. On administration of the naproxen formulations of the present invention, the formulation, preferably a tablet, rapidly dissociates releasing the pellets over a wide area of the GI  
20       tract. Thus, the number of gastrointestinal absorption sites is increased minimizing the occurrence of adverse GI effects often associated with the administration of NSAIDs such as naproxen. The subject naproxen tablet formulation therefore is ideally suited to a once-daily dosage regimen having the potential for fewer associated adverse GI effects.

- 25       The accompanying Figure is a graph of mean plasma levels ( $\mu\text{g/ml}$ ) versus time (hours) for a naproxen formulation according to the invention relative to two reference naproxen formulations.

The invention will be further illustrated by the following Examples:

30

12

2034096

Example 1

Naproxen sodium (50 kg), citric acid (14.286 kg), sodium lauryl sulphate (1.071 kg) and talc (3.571 kg) were blended and milled through a suitable mesh screen so as to obtain a homogenous powder.

- 5        The powder was applied to starch/sugar seeds (0.3 - 0.425 mm diameter) (4.286 kg) using a FREUND CF granulator and a coating solution of 3.5% polyvinylpyrrolidone in isopropanol to form the cores.

A membrane was then applied to the cores by spraying on a solution consisting of:

10	12.5% EUDRAGIT RS in acetone/isopropanol 40:60	50 parts by weight
	12.5% EUDRAGIT RL in acetone/isopropanol 40:60	40 parts by weight
15	12.5% EUDRAGIT L in acetone/isopropanol 40:60	10 parts by weight
	Isopropanol	100 parts by weight

- while at the same time but separately dusting on talc (100 parts by weight) in conventional manner. The ratio of membrane solution to talc applied was 0.62 g of talc per gram of membrane solution. A sufficient  
 20        amount of membrane solution and talc was applied to 76.214 kg of cores to achieve the dissolution profile given below.

The finished pellets were dried to evaporate all solvents prior to performing the dissolution profile.

- 25        A wet granulation of naproxen sodium (29.100 kg) and polyvinylpyrrolidone (K30) (0.900 kg) was prepared by adding isopropanol (8.2351 kg) slowly with mixing. Mixing was continued until a uniform mass was produced. The granulation was dried for a

13

2034096

minimum of 12 hours at a temperature of not less than 45°C. The dried granulation was then passed through an oscillating granulator.

Naproxen sodium pellets (27.216 kg) and naproxen granulate (8.112 kg) were blended with additional tableting excipients:

5	Microcrystalline cellulose	15.372 kg
	Crosspovidone	1.049 kg
	Magnesium stearate	0.260 kg

and pressed into uncoated tablets.

10 Slight adjustments may be made in the above quantities to ensure a target tablet potency of 500 mg of naproxen.

15 The naproxen tablets were coated with an aqueous film coat. To do so, 0.350 kg of Opadry White OY-D-7192 (Opadry White is a Trade Mark) was weighed and slowly added with mixing to 1.400 kg of purified water. Mixing was continued for 30 minutes. 10 kg of uncoated tablets were weighed and placed in an ACCELA-COTA. The coating was sprayed onto the tablets at a rate of 30-70 g per minute until coating was complete.

20 The dissolution rate of the pellets prepared above, prior to mixing with the granulate, was tested by the method of the U.S. Pharmacopoeia XXII Basket Method in phosphate buffer at pH 7.2 and at 75 r.p.m. The naproxen sodium was quantitatively determined using u.v. spectrophotometry at 331 nm.

25

14

2034096

The dissolution rate was as follows:

	<u>Time (h)</u>	<u>% Naproxen Sodium Released</u>
5	1	11.2
	2	42.6
	4	80.3

The dissolution rate of the coated tablets was tested and determined to be as follows:

	<u>Time (h)</u>	<u>% Naproxen Sodium Released</u>
10	0.5	36.8
	1	50.5
	4	95.3

#### Example 2

15 Naproxen sodium (15.4 kg), citric acid (4.4 kg) and talc (0.200kg) were blended and milled through a suitable mesh screen to obtain a homogeneous powder. This powder blend was layered into spherical cores using a FREUND CF granulator or a GLATT and a coating solution of 12.5% Eudragit RS in isopropanol.

20 A membrane was then applied to the cores by spraying on a solution consisting of:

	12.5% EUDRAGIT RS in acetone/isopropanol 40:60	70 parts by weight
25	12.5% EUDRAGIT RL in acetone/isopropanol 40:60	10 parts by weight



15

2034096

12.5% EUDRAGIT L in  
acetone/isopropanol 40:60 20 parts by weight

Isopropanol 100 parts by weight

5 while at the same time but separately dusting on talc (40 parts by weight) in the conventional manner. The ratio of membrane solution to talc applied was 0.25 g of talc per gram of membrane solution. A sufficient amount of membrane solution and talc was applied to 20.00 kg of cores to achieve the dissolution profile given below.

10 The finished pellets were dried to evaporate all solvents prior to performing the dissolution test.

A wet granulation of naproxen sodium (9.700 kg) and polyvinylpyrrolidone (K30) (0.300 kg) was prepared by adding isopropanol (2.745 kg) slowly with mixing. Mixing was continued until a uniform mass was produced. The granulation was dried for a  
15 minimum of 12 hours at a temperature of not less than 45°C. The dried granulation was then passed through an oscillating granulator.

Naproxen sodium pellets (24 kg) and naproxen granulate (6.4 kg) were blended with additional tableting excipients:

20	Microcrystalline cellulose	4.63 kg
	Crosspovidone	0.76 kg
	Magnesium stearate	0.19 kg
	PVP K30	0.76 kg

and pressed into uncoated tablets.

25 Slight adjustments may be made in the above quantities to ensure a target potency of 500 mg of naproxen.

The naproxen tablets were coated with an aqueous film coat. To do so, 0.350 kg of Opadry White OY-D-7192 was weighed and slowly

16

2034096

added with mixing to 1.400 kg of purified water. Mixing was continued for 30 minutes. 10 kg of uncoated tablets were weighed and placed in an ACCELA-COTA. The coating was sprayed onto the tablets at a rate of 30-70 g per minute until coating was complete.

- 5        The dissolution rate of the pellets prepared above, prior to mixing with the granulate, was tested by the method of the U.S. Pharmacopoeia XXII Basket Method in phosphate buffer at pH 7.2 and at 75 r.p.m. The naproxen sodium was quantitatively determined using u.v. spectrophotometry at 331 nm.

- 10       The dissolution rate was as follows:

	<u>Time (h)</u>	<u>% Naproxen Sodium Released</u>
	1	23.9
	2	36.6
15	4	61.3

The dissolution rate of the coated tablets was tested and determined to be as follows:

	<u>Time (h)</u>	<u>% Naproxen Sodium Released</u>
20	1	60.0
	2	78.3
	4	89.2

### Example 3

- 25       Naproxen sodium (25 kg), citric acid (7.141 kg) and talc (0.325 kg) were blended and milled through a suitable mesh screen so as to obtain a homogeneous powder. This powder blend was divided into two portions and layered into spherical cores using a FREUND CP

17

2034096

granulator or a GLATT and a coating solution of 12.5% Eudragit RS in isopropanol.

A membrane was then applied to the cores by spraying on a solution consisting of:

5	12.5% EUDRAGIT RS in acetone/isopropanol 40:60	75 parts by weight
	12.5% EUDRAGIT RL in acetone/isopropanol 40:60	25 parts by weight
	Isopropanol	100 parts by weight

10 while at the same time but separately dusting on talc (10 parts by weight) in the conventional manner. The ratio of membrane solution to talc applied was 0.06 g of talc per gram of membrane solution. A sufficient amount of membrane solution and talc was applied to 33.4 kg of cores to achieve the dissolution profile given below.

15 The finished pellets were dried to evaporate all solvents prior to performing the dissolution test.

20 A wet granulation of naproxen sodium (19.400 kg) and polyvinylpyrrolidone (K30) (0.600 kg) was prepared by adding isopropanol (1.529 kg) slowly with mixing. Mixing was continued until a uniform mass was produced. The granulation was dried for a minimum of 12 hours at a temperature of not less than 45°C. The dried granulation was then passed through an oscillating granulator.

Naproxen sodium pellets (25 kg) and naproxen granulate (7.85 kg) were blended with additional tableting excipients:

25	Microcrystalline cellulose	6.00 kg
	Crosspovidone	0.815 kg

18

2034096

Magnesium stearate	0.200 kg
PVP K30	0.815 kg

and pressed into uncoated tablets.

5 Slight adjustments may be made in the above quantities to ensure a target tablet potency of 500 mg of naproxen.

10 The naproxen tablets were coated with an aqueous film coat. To do so, 0.350 kg of Opadry White OY 9400 was weighed and slowly added with mixing to 1.400 kg of purified water. Mixing was continued for 30 minutes. 10 kg of uncoated tablets were weighed and placed in an ACCBLA-COTA. The coating was sprayed onto the tablets at a rate of 30-70 g per minute until coating was complete.

15 The dissolution rate of the pellets prepared above, prior to mixing with the granulate, was tested by the method of the U.S. Pharmacopoeia XXII Basket Method in phosphate buffer at pH 7.2 and at 75 r.p.m. The naproxen sodium was quantitatively determined using u.v. spectrophotometry at 331 nm.

The dissolution rate was as follows:

	<u>Time (h)</u>	<u>% Naproxen Sodium Released</u>
20	1	30.1
	2	42.5
	4	61.5

The dissolution rate of the coated tablets was tested and determined to be as follows:

25

2034096

19

	<u>Time (h)</u>	<u>% Naproxen Sodium Released</u>
	0.5	39.6
	1	55.0
5	4	90.0

Example 4

Naproxen acid (50 kg), citric acid (15.64 kg), sodium lauryl sulphate (0.071 kg) and talc (0.500 kg) were blended and milled through a suitable mesh screen to obtain a homogeneous powder. This powder blend was layered into spherical cores using a FREUND CF granulator or a GLATT and a coating solution of 12.5% Eudragit RS in isopropanol.

A membrane was then applied to the cores by spraying on a solution consisting of:

15	12.5% EUDRAGIT RS in acetone/isopropanol 40:60	90 parts by weight
	12.5% EUDRAGIT RL in acetone/isopropanol 40:60	10 parts by weight
	Isopropanol	100 parts by weight

while at the same time but separately dusting on talc (20 parts by weight) in the conventional manner. The ratio of membrane solution to talc applied was 0.125 g of talc per gram of membrane solution. A sufficient amount of membrane solution and talc was applied to 66.22 kg of cores to achieve the dissolution profile given below.

The finished pellets were dried to evaporate all solvents prior to performing the dissolution test. The dissolution rate of the pellets was

20

2034096

tested by the method of the U.S. Pharmacopoeia XXI: Basket Method in phosphate buffer, at pH 7.2 and at 75 r.p.m.

The naproxen was quantitatively determined using u.v. spectrophotometry at 331 nm.

5        The dissolution rate was as follows:

	<u>Time (h)</u>	<u>% Naproxen Released</u>
	1	23.9
	2	36.6
10	4	61.3

Example 5

Naproxen cores were prepared according to Example 1. A membrane was then applied to the cores by spraying on a suspension of:

	30% BUDRAGIT RS in water	30 parts by weight
15	30% BUDRAGIT RL in water	6 parts by weight
	Talc	6 parts by weight
	Water	50 parts by weight

The remaining 8 parts by weight were made up by a 9:1 mixture of tributylacetyl citrate, a plasticizer, and simethicone, an antifoam agent.

20    A sufficient amount of membrane solution was applied to achieve the following desired release rate.

21

2034096

	<u>Time (h)</u>	<u>% Naproxen Sodium Released</u>
	1	29.8
	2	47.7
5	4	73.6

Example 6

Naproxen cores were prepared according to Example 3.

A membrane was then applied to the cores by spraying on a suspension of:

10	30% EUDRAGIT <sup>®</sup> RS in water	40 parts by weight
	30% EUDRAGIT <sup>®</sup> RL in water	2 parts by weight
	Talc	6 parts by weight
	Water	50 parts by weight

15 The remaining 2 parts by weight were made up by a 9:1 mixture of tributylacetyl citrate, a plasticizer, and simethicone, an antifoam agent.

A sufficient amount of membrane solution was applied to achieve the desired release rate of:

	<u>Time (h)</u>	<u>% Naproxen Sodium Released</u>
20	1	14.5
	2	42.7
	4	71.1

Example 7

25 Naproxen sodium (30.0 kg), citric acid (5.00 kg) and talc (0.400 kg) were blended and milled through a suitable mesh screen to obtain a

homogeneous powder. This powder blend was layered into spherical cores using a PRBUND CP granulator or a GLATT and a coating solution of 12.5% Eudragit RS in isopropanol.

- 5 A membrane was then applied to the cores by spraying on a solution consisting of:

	12.5% EUDRAGIT RS in acetone/isopropanol 40:60	85 parts by weight
	12.5% EUDRAGIT RL in acetone/isopropanol 40:60	15 parts by weight
10	Isopropanol	100 parts by weight

- while at the same time but separately dusting on talc (40 parts by weight) in the conventional manner. The ratio of membrane solution to talc applied was 0.25 g of talc per gram of membrane solution. A sufficient amount of membrane solution and talc was applied to 35.4 kg of cores to achieve the dissolution profile given below.

The finished pellets were dried to evaporate all solvents prior to performing the dissolution test.

- 20 A wet granulation of naproxen sodium (14.550 kg) and polyvinylpyrrolidone (K30) (0.450 kg) was prepared by adding isopropanol (4.116 kg) slowly with mixing. Mixing was continued until a uniform mass was produced. The granulation was dried for a minimum of 12 hours at a temperature of not less than 45°C. The dried granulation was then passed through an oscillating granulator.

- 25 Naproxen sodium pellets (35.0 kg) and naproxen granulate (8.10 kg) were blended with additional tableting excipients:

Microcrystalline cellulose	8.346 kg
Crosspovidone	1.077 kg



23

2034036

Magnesium stearate	0.270 kg
PVP K30	1.077 kg

and pressed into uncoated tablets.

5      Slight adjustments may be made in the above quantities to ensure  
a target tablet potency of 750 mg of naproxen.

The naproxen tablets were coated with an aqueous film coat. To  
do so, 0.350 kg of Opadry White OY 9400 was weighed and slowly  
added with mixing to 1.400 kg of purified water. Mixing was continued  
for 30 minutes. 10 kg of uncoated tablets were weighed and placed in an  
10      ACCELA-COTA. The coating was sprayed onto the tablets at a rate of  
30-70 g per minute until coating was complete.

The dissolution rate of the pellets prepared above, prior to  
mixing with the granulate, was tested by the method of the U.S.  
Pharmacopocia XXII Basket Method in phosphate buffer at pH 7.2 and  
15      at 75 r.p.m. The naproxen sodium was quantitatively determined using  
u.v. spectrophotometry at 331 nm.

The dissolution rate was as follows:

	<u>Time (h)</u>	<u>% Naproxen Sodium Released</u>
20	1	50.7
	2	71.3
	4	88.0

The dissolution rate of the coated tablets was tested and  
determined to be as follows:

25

24

2034096

	<u>Time (h)</u>	<u>% Naproxen Sodium Released</u>
	0.5	49.5
	1	71.8
5	4	89.4

Example 8

Naproxen sodium (22.5 kg), citric acid (1.125 kg) and talc (0.251 kg) were blended and milled through a suitable mesh screen to obtain a homogeneous powder. This powder blend was layered into spherical  
 10 cores using a FREUND CP granulator or a GLATT and a coating solution of 12.5% Eudragit RS in isopropanol.

A membrane was then applied to the cores by spraying on a solution consisting of:

15	12.5% EUDRAGIT RS in acetone/isopropanol 40:60	70 parts by weight
	12.5% EUDRAGIT RL in acetone/isopropanol 40:60	10 parts by weight
	12.5% EUDRAGIT L in acetone/isopropanol 40:60	20 parts by weight
20	Isopropanol	100 parts by weight

while at the same time but separately dusting on talc (5 part by weight) in the conventional manner. The ratio of membrane solution to talc applied was 0.03 g of talc per gram of membrane solution. A sufficient  
 25 amount of membrane solution and talc was applied to 12.48 kg of cores to achieve the dissolution profile given below.

25

2034096

The finished pellets were dried to evaporate all solvents prior to performing the dissolution test.

5 A wet granulation of naproxen sodium (14.55 kg) and polyvinylpyrrolidone (K30) (0.45 kg) was prepared by adding isopropanol (1.147 kg) slowly with mixing. Mixing was continued until a uniform mass was produced. The granulation was dried for a minimum of 12 hours at a temperature of not less than 45°C. The dried granulation was then passed through an oscillating granulator.

10 Naproxen sodium pellets (12.0 kg) and naproxen granulate (3.986 kg) were blended with additional tableting excipients:

Microcrystalline cellulose	2.32 kg
Crosspovidone	0.58 kg
Magnesium stearate	0.10 kg
PVP K30	0.38 kg

15 and pressed into uncoated tablets.

Slight adjustments may be made in the above quantities to ensure a target tablet potency of 750 mg of naproxen.

20 The naproxen tablets were coated with an aqueous film coat. To do so, 0.350 kg of Opadry White OY-D-7192 was weighed and slowly added with mixing to 1.400 kg of purified water. Mixing was continued for 30 minutes. 10 kg of uncoated tablets were weighed and placed in an ACCBLA-COTA. The coating was sprayed onto the tablets at a rate of 30-70 g per minute until coating was complete.

25 The dissolution rate of the pellets prepared above, prior to mixing with the granulate, was tested by the method of the U.S. Pharmacopoeia XXII Basket Method in phosphate buffer at pH 7.2 and at 75 r.p.m. The naproxen sodium was quantitatively determined using u.v. spectrophotometry at 331 nm.

26

2034088

The dissolution rate was as follows:

	<u>Time (h)</u>	<u>% Naproxen Sodium Released</u>
5	1	23.9
	2	36.6
	4	61.4

The dissolution rate of the coated tablets was tested and determined to be as follows:

	<u>Time (h)</u>	<u>% Naproxen Sodium Released</u>
10	0.5	40.8
	1	70.0
	4	87.0

\* \* \* \* \*

#### 15 Pharmacological Data for Naproxen Formulations

##### In Vivo Performance:

##### Pharmacological Data for Naproxen Formulation of Example 1

The naproxen formulation prepared in Example 1 was evaluated *in vivo* under steady state conditions.

- 20 A steady state study was performed in 18 healthy male volunteers, comparing the formulation of Example 1 with selected reference products, i.e., conventional immediate release tablets. The formulation of Example 1 was administered as a single tableted dose of 1,000 mg of naproxen at 0 hours on days 1-7, while reference Naprosyn (N), a
- 25 commercially available naproxen formulation manufactured by Syntex which is administered twice daily, was administered as a single 500 mg

27

2034096

tablet at 0 and 12 hours (i.e. b.i.d.) on days 1-7 and reference Syntex  
 (S), a commercially available naproxen sodium formulation  
 manufactured by Syntex which is administered three to four times daily,  
 was administered as a single 250 mg dose at 0, 6, 12 and 18 hours (i.e.  
 5 q.i.d.) on days 1-7. Plasma was sampled out to 24 hours and the mean  
 results were calculated and tabulated. The results are illustrated in the  
 accompanying Figure. In the latter Figure curve a) corresponds to a  
 naproxen formulation prepared according to Example 1; curve b)  
 corresponds to Reference (N); and curve c) corresponds to Reference  
 10 (S).

The data presented in Table 1 are from day 7 sampling.

Table 1

Mean Naproxen Concentrations ( $\mu\text{g/ml}$ ) - Day 7

<u>Hour</u>	<u>Reference (N)</u>	<u>Reference (S)</u>	<u>Formulation of Example 1</u>
0.0	46.95	59.80	42.39
0.50	86.41	84.92	74.95
1.00	102.09	94.33	84.89
2.00	99.33	77.63	90.15
4.00	76.38	65.05	93.09
6.00	63.34	53.90	91.22
6.50	59.20	58.21	87.26
7.00	57.10	68.53	88.48
8.00	52.76	72.98	80.64
10.00	48.56	69.52	69.02
12.00	41.16	57.07	65.68
12.50	55.27	62.46	63.52
13.00	67.12	72.45	60.83
14.00	83.94	74.81	57.78
16.00	80.05	67.10	51.26
18.00	68.39	55.96	44.37
18.50	63.09	66.89	45.84
19.00	62.14	74.40	44.63
20.00	58.99	72.92	42.89
22.00	53.39	68.11	39.97
24.00	47.90	61.38	38.30

The results of this *in vivo* comparison of the formulation of Example 1 against conventional immediate release tablets (Reference N and S) indicate the formulation of Example 1 to be bioequivalent (99.28%) to Reference N and bioequivalent (95.46%) to Reference S. Both Example 1 and Reference S formulations use the sodium salt of naproxen, the Reference N formulation, however, uses the free acid form of naproxen. Generally, peak plasma concentrations following administration of naproxen as its sodium salt are higher and are attained more rapidly than those following administration of naproxen as the free acid, hence the use of the sodium salt in the conventional form as an analgesic on a more frequent dosing regimen (3 - 4 times daily).

The  $T_{max}$  value of the formulation of the present invention of 5.0 hours was significantly longer than that of Reference N at 1.36 hours, thus demonstrating the improved suitability of the present formulations in terms of maintaining naproxen plasma levels important for continued analgesia and anti-inflammatory effects up to 24 hours following administration. Furthermore, the formulation of Example 1 exhibited an extended mean plasma profile with adequate duration of time cover in the range of desired therapeutic plasma naproxen concentrations of 30-60  $\mu\text{g/ml}$ . Also, the mean  $C_{max}$  value of the product of Example 1 was close to the  $C_{max}$  values of the References, indicating that the sustained release characteristics of Example 1 did not adversely affect the maximum plasma levels attained following administration of the formulation.

A gastric tolerability questionnaire was completed by each subject on each day of the study. The results indicated a 47% reduction in adverse gastrointestinal effects for the formulation prepared according to Example 1 when compared to those recorded by Reference N or S.

The clinical efficacy of the naproxen formulation according to the present invention was evaluated in osteoarthritic patients. In osteoarthritis, NSAIDs are primarily used for their analgesic rather

than their anti-inflammatory effect, although inflammation may be part of the symptomatology. To evaluate the formulation of the present invention, the formulation of Example 1 was compared with Naprosyn in a double-blind double placebo controlled study in osteoarthritic patients.

Efficacy of both naproxen formulations was indicated by the degree of pain assessed by the patient on a visual analogue scale and was graded on a four point verbal rating scale. In addition, blood samples were analysed for naproxen plasma concentrations and any adverse effects were recorded. A gastrointestinal tolerability questionnaire was completed by each volunteer at the end of each treatment period.

In summary, in terms of clinical efficacy, the naproxen formulation according to the present invention performed as well as the reference and was superior to the reference in terms of fewer gastrointestinal complaints and adverse events, thus demonstrating the safety and advantage of the multiparticulate formulations of the present invention when compared with the commercially available reference products.

CA 02034096 2001-01-19

30

CLAIMS:

1. A naproxen formulation for oral administration on a once-daily basis, said formulation comprising a first portion of the naproxen formulated in a multi-particulate pellet form, each pellet having a core  
5 of naproxen or a pharmaceutically acceptable salt thereof surrounded by a multi-layer membrane, said membrane containing a major proportion of a pharmaceutically acceptable film-forming, water insoluble polymer and optionally a minor proportion of a pharmaceutically acceptable film-forming, water soluble polymer,  
10 said multi-particulate form being effective to release the drug at such a rate as to maintain therapeutically effective blood levels substantially over 24 hours when administered each day on a once-daily basis and, optionally, a second portion of said naproxen formulated so as to release the drug promptly following  
15 administration so as to obtain a relatively immediate therapeutic response.
2. A naproxen formulation according to claim 1, wherein said first portion is in pellet form, said pellet comprising a core of naproxen or a pharmaceutically acceptable salt thereof in association with an  
20 organic acid, the naproxen component and the organic acid being present in a ratio of from 20:1 to 1:1, and a multi-layer membrane surrounding said core and containing a major proportion of a pharmaceutically acceptable film-forming, water insoluble polymer and optionally a minor proportion of a pharmaceutically acceptable  
25 film-forming, water soluble polymer, the number of layers in said membrane and the ratio of said water soluble to water insoluble polymer being effective to permit release of said naproxen from said



CA 02034096 2001-01-19

30 a

pellet at a rate allowing controlled absorption thereof at therapeutically effective blood levels over a 24 hour period following each daily oral administration.

3. A pellet formulation according to Claim 2, wherein the pellet  
5 has a dissolution rate which when measured *in vitro* in a type 1 dissolution basket apparatus according to U.S. Pharmacopoeia XXII in phosphate buffer at pH 7.2 and at 75 r.p.m. substantially corresponds to the following dissolution pattern:

- 10 a) from 0 to 50% of the total naproxen is released after 1 hour of measurement in said apparatus;
- b) from 20 to 70% of the total naproxen is released after 2 hours of measurement in said apparatus; and

- c) not less than 50% of the total naproxen is released after 4 hours of measurement in said apparatus.
4. A pellet formulation according to Claim 2, wherein the pellet has a dissolution rate which when measured *in vitro* in a type 2 dissolution basket apparatus according to U.S. Pharmacopoeia XXII in phosphate buffer at pH 7.4 and at 50 r.p.m. the dissolution rate substantially corresponds to the following dissolution pattern:
- 5
- a) from 20 to 70% of the total naproxen is released after 1 hour of measurement in said apparatus;
- 10
- b) not less than 50% of the total naproxen is released after 2 hours of measurement in said apparatus; and
- c) not less than 75% of the total naproxen is released after 4 hours of measurement in said apparatus.
5. A pellet formulation according to Claim 2, wherein the organic acid is selected from one or more of the following acids: adipic acid, ascorbic acid, citric acid, fumaric acid, malic acid, succinic acid or tartaric acid.
- 15
6. A pellet formulation according to Claim 2, wherein the naproxen or pharmaceutically acceptable salt thereof and organic acid are present in a ratio of from 20:1 to 1:2.
- 20
7. A pellet formulation according to Claim 2, wherein the naproxen or pharmaceutically acceptable salt thereof and organic acid are present in a ratio of 6:1 to 1:1.
8. A pellet formulation according to Claim 2, wherein the core includes a lubricant selected from one or more of the following: sodium stearate, magnesium stearate, stearic acid or talc.
- 25

9. A pellet formulation according to Claim 2, wherein the core includes a lubricant selected from one or more of the following: sodium stearate, magnesium stearate, stearic acid or talc and the naproxen and lubricant are present in a ratio of from 10:1 to 200:1.
- 5 10. A pellet formulation according to Claim 2, wherein the core comprises naproxen or a pharmaceutically acceptable salt thereof and the associated organic acid embedded in a polymeric material.
11. A pellet formulation according to Claim 2, wherein the core comprises naproxen or a pharmaceutically acceptable salt thereof and  
10 the associated organic acid embedded in a polymeric material, said polymeric material being insoluble in water or slightly permeable to naproxen and water.
12. A pellet formulation according to Claim 2, wherein the core comprises naproxen or a pharmaceutically acceptable salt thereof and  
15 the associated organic acid embedded in a polymeric material, said polymeric material consisting solely of a water insoluble polymer or a polymer which is slightly permeable to water and aqueous solutions of naproxen.
13. A pellet formulation according to Claim 2, wherein the core  
20 comprises naproxen or a pharmaceutically acceptable salt thereof and the associated organic acid embedded in a polymeric material, said polymeric material consisting solely of a water soluble polymer or a polymer which is freely permeable to aqueous solutions of naproxen and water.
14. A pellet formulation according to Claim 2, wherein the core  
25 comprises naproxen or a pharmaceutically acceptable salt thereof and the associated organic acid embedded in a polymeric material, said polymeric material comprising a combination of a water insoluble polymer with a water soluble polymer, the ratio of water soluble/freely  
30 permeable to water insoluble/slightly permeable polymer being determined by the particular combination of polymers selected.

15. A pellet formulation according to Claim 2, wherein the core comprises:
- 5 a) a powder mixture containing naproxen or a pharmaceutically acceptable salt thereof, an organic acid selected from adipic acid, ascorbic acid, citric acid, fumaric acid, malic acid, succinic acid and tartaric acid; and
- 10 b) a polymeric material containing a proportion of a pharmaceutically acceptable water soluble polymer and a proportion of a pharmaceutically acceptable water insoluble polymer, said core comprising layers of said powder mixture and said polymeric material superimposed one upon the other and said polymeric material being present in an amount effective to ensure that all of said powder mixture is coated into said core.
- 15 16. A pellet formulation according to Claim 2, wherein the core comprises:
- a) a powder mixture containing naproxen or a pharmaceutically acceptable salt thereof, an organic acid selected from adipic acid, ascorbic acid, citric acid, fumaric acid, malic acid, succinic acid and tartaric acid; and
- 20 b) a polymeric material containing a proportion of a pharmaceutically acceptable water soluble polymer and a proportion of a pharmaceutically acceptable water insoluble polymer, said core comprising layers of said powder mixture and said polymeric material superimposed one upon the other and said
- 25 polymeric material being present in an amount effective to ensure that all of said powder mixture is coated into said core.

and wherein the water soluble polymer is selected from polyvinyl alcohol, polyvinylpyrrolidone, methyl cellulose, hydroxypropylcellulose, agar, carrageenan, xanthan,

hydroxypropylmethyl cellulose or polyethylene glycol or a mixture thereof.

17. A pellet formulation according to Claim 2, wherein the core comprises:

- 5 a) a powder mixture containing naproxen or a pharmaceutically acceptable salt thereof, an organic acid selected from adipic acid, ascorbic acid, citric acid, fumaric acid, malic acid, succinic acid and tartaric acid; and
- 10 b) a polymeric material containing a proportion of a pharmaceutically acceptable water soluble polymer and a proportion of a pharmaceutically acceptable water insoluble polymer, said core comprising layers of said powder mixture and said polymeric material superimposed one upon the other and said polymeric material being present in an amount effective to ensure
- 15 that all of said powder mixture is coated into said core,

and wherein the polymeric material is freely permeable to naproxen and water and comprises a copolymer of acrylic and methacrylic acid esters.

18. A pellet formulation according to Claim 2, wherein the core comprises:

- 20 a) a powder mixture containing naproxen or a pharmaceutically acceptable salt thereof, an organic acid selected from adipic acid, ascorbic acid, citric acid, fumaric acid, malic acid, succinic acid and tartaric acid; and
- 25 b) a polymeric material containing a proportion of a pharmaceutically acceptable water soluble polymer and a proportion of a pharmaceutically acceptable water insoluble polymer, said core comprising layers of said powder mixture and said polymeric material superimposed one upon the other and said

35

2034096

polymeric material being present in an amount effective to ensure that all of said powder mixture is coated into said core,

and wherein the water insoluble polymer is selected from ethyl cellulose, cellulose acetate, cellulose propionate (lower, medium or higher molecular weight), cellulose acetate propionate, cellulose acetate butyrate, cellulose acetate phthalate, cellulose triacetate, poly(methyl methacrylate), poly(ethyl methacrylate), poly(butyl methacrylate), poly(isobutyl methacrylate), poly(hexyl methacrylate), poly(isodecyl methacrylate), poly(lauryl methacrylate), poly(phenyl methacrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), poly(octadecyl acrylate), poly(ethylene), poly(ethylene) low density, poly(ethylene) high density, poly(propylene), poly(ethylene oxide), poly(ethylene terephthalate), poly(vinyl isobutyl ether), poly(vinyl acetate), poly(vinyl chloride) or polyurethane or a mixture thereof.

19. A pellet formulation according to Claim 2, wherein the core comprises:

- a) a powder mixture containing naproxen or a pharmaceutically acceptable salt thereof, an organic acid selected from adipic acid, ascorbic acid, citric acid, fumaric acid, malic acid, succinic acid and tartaric acid; and
- b) a polymeric material containing a proportion of a pharmaceutically acceptable water soluble polymer and a proportion of a pharmaceutically acceptable water insoluble polymer, said core comprising layers of said powder mixture and said polymeric material superimposed one upon the other and said polymeric material being present in an amount effective to ensure that all of said powder mixture is coated into said core,

and wherein the water insoluble polymer is selected from a naturally occurring polymer selected from shellac, chitosan, gumjuniper or a mixture thereof.

36

2034096

20. A pellet formulation according to Claim 2, wherein the core comprises:

- 5 a) a powder mixture containing naproxen or a pharmaceutically acceptable salt thereof, an organic acid selected from adipic acid, ascorbic acid, citric acid, fumaric acid, malic acid, succinic acid and tartaric acid; and
- 10 b) a polymeric material containing a proportion of a pharmaceutically acceptable water soluble polymer and a proportion of a pharmaceutically acceptable water insoluble polymer, said core comprising layers of said powder mixture and said polymeric material superimposed one upon the other and said polymeric material being present in an amount effective to ensure that all of said powder mixture is coated into said core,

15 and wherein the polymeric material is slightly permeable to naproxen and water and comprises a copolymer of acrylic and methacrylic acid esters.

21. A pellet formulation according to Claim 2, wherein the core comprises naproxen or a pharmaceutically acceptable salt thereof and the associated organic acid embedded in a polymeric material, and  
20 wherein the naproxen, organic acid and polymeric material are built up on a central active core.

22. A pellet formulation according to Claim 2, wherein the water insoluble polymer of the membrane is selected from ethyl cellulose, cellulose acetate, cellulose propionate (lower, medium or higher  
25 molecular weight), cellulose acetate propionate, cellulose acetate butyrate, cellulose acetate phthalate, cellulose triacetate, poly(methyl methacrylate), poly(ethyl methacrylate), poly(butyl methacrylate), poly(isobutyl methacrylate), poly(hexyl methacrylate), poly(isodecyl methacrylate), poly(lauryl methacrylate), poly(phenyl methacrylate),  
30 poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), poly(octadecyl acrylate), poly(ethylene), poly(ethylene) low density,

poly(ethylene) high density, poly(propylene), poly(ethylene oxide), poly(ethylene terephthalate), poly(vinyl isobutyl ether), poly(vinyl acetate), poly(vinyl chloride) or polyurethane or a mixture thereof.

23. A pellet formulation according to Claim 2, wherein the water  
5 insoluble polymer of the membrane is selected from a naturally occurring polymer selected from shellac, chitosan, gumjuniper or a mixture thereof.

24. A pellet formulation according to Claim 2, wherein the water  
soluble polymer of the membrane is selected from polyvinyl alcohol,  
10 polyvinylpyrrolidone, methyl cellulose, hydroxypropyl cellulose, agar, carrageenan, xanthan, hydroxypropylmethyl cellulose or polyethylene glycol or a mixture thereof.

25. A pellet formulation according to Claim 2, wherein the water  
insoluble polymer of the membrane is replaced as appropriate by a  
15 proportion of a polymer which is slightly permeable to naproxen and water and the water soluble polymer of the membrane is replaced by a proportion of a polymer which is freely permeable to naproxen and water.

26. A pellet formulation according to Claim 2, wherein the water  
20 insoluble polymer of the membrane is replaced as appropriate by a proportion of a polymer which is slightly permeable to naproxen and water and the water soluble polymer of the membrane is replaced by a proportion of a polymer which is freely permeable to naproxen and water, and wherein each of the slightly permeable and freely permeable  
25 polymers comprises a copolymer of acrylic and methacrylic acid esters having the desired permeability characteristics.

27. A pellet formulation according to Claim 2, which includes in the  
core and/or the membrane, a plasticizing agent selected from  
polyethylene glycol, propylene glycol, glycerol, triacetin, dimethyl  
30 phthalate, diethyl phthalate, dibutyl phthalate, dibutyl sebacate, triethyl



CA 02034096 2001-01-19

38

citrate, tributyl citrate, triethyl acetyl citrate, castor oil or an acetylated monoglyceride.

28. A formulation according to Claim 2, which contains naproxen sodium as the active ingredient.

5 29. A process for the production of a formulation according to Claim 2, which comprises forming a core of naproxen or a pharmaceutically acceptable salt thereof and an organic acid and enclosing the core in a membrane of a film-forming polymer or mixture thereof as defined in Claim 2 which permits release of the naproxen or the pharmaceutically  
10 acceptable salt thereof in the manner set out in Claim 2.

30. A controlled absorption naproxen formulation for oral administration, comprising pellets each pellet comprising a core of naproxen or a pharmaceutically acceptable salt thereof in association with an organic acid, the naproxen component and the organic acid  
15 being present in a ratio of from 20:1 to 1:1, and a multi-layer membrane surrounding said core and containing a major proportion of a pharmaceutically acceptable film-forming, water insoluble polymer and optionally a minor proportion of a pharmaceutically acceptable film-forming, water soluble polymer, the number of layers in said  
20 membrane and the ratio of said water soluble to water insoluble polymer being effective to permit release of said naproxen from said pellet at a rate allowing controlled absorption thereof at therapeutically effective blood levels over a 24 hour period following each daily oral administration, said formulation including a sufficient quantity of a  
25 rapid release form of naproxen to ensure prompt achievement of analgesically effective blood levels together with therapeutically effective blood levels over a 24 hour period following each daily oral administration.

31. A controlled absorption naproxen formulation according to Claim  
30 30, which has a dissolution rate which when measured in a type 1 dissolution basket apparatus according to U.S. Pharmacopoeia XXII in

39

2034096

phosphate buffer at pH 7.2 and at 75 r.p.m. substantially corresponds to the following dissolution pattern:

- a) from 15 to 50% of the total naproxen is released after 0.5 hours of measurement in said apparatus;
- 5 b) from 25 to 75% of the total naproxen is released after 1 hour of measurement in said apparatus; and
- c) not less than 65% of the total naproxen is released after 4 hours of measurement in said apparatus.

32. A controlled absorption naproxen formulation according to Claim 10 30, which has a dissolution rate which when measured *in vitro* in a type 2 dissolution paddle apparatus according to U.S. Pharmacopoeia XXII in phosphate buffer at pH 7.4 and at 50 r.p.m. the dissolution rate substantially corresponds to the following dissolution pattern:

- 15 a) from 25 to 60% of the total naproxen is released after 0.5 hours of measurement in said apparatus;
- b) from 35 to 75% of the total naproxen is released after 1 hour of measurement in said apparatus; and
- c) not less than 65% of the total naproxen is released after 4 hours of measurement in said apparatus.

20 33. A controlled absorption naproxen formulation according to Claim 30, which comprises a blend of said pellets, together with up to 60% by weight of said rapid release form of naproxen.

25 34. A controlled absorption naproxen formulation according to Claim 30, wherein the rapid release form of naproxen comprises a rapid release granulate.

40

2034096

35. A formulation according to Claim 1, 2 or 30 in the form of a capsule or tablet.


2034096

ABSTRACT

CONTROLLED ABSORPTION NAPROXEN FORMULATION FOR  
ONCE-DAILY ADMINISTRATION

A naproxen formulation for oral administration on a once-daily basis comprises a first portion of the naproxen formulated in a multi-particulate pellet form designed to release the drug at a rate sufficient to maintain therapeutically effective blood levels substantially over 24 hours when administered each day on a once-daily basis and, optionally, a second portion of the naproxen formulated so as to release the drug promptly following administration so as to obtain a relatively immediate therapeutic response.

2034096



Spicy, India, Galt  
Baker & Brown